

P.O. Box 280173, Tampa, Florida 33682 (813) 988-0115

## ERRATA

In the January-March, 1988 issue, the proceedings for the Protocol Roundtable on Acute Congestive Heart Failure (*Protocol Roundtable: Acute Congestive Heart Failure. Tampa Bay EMS Journal 1(3):62-70, 1988*) contains errors in Figure 6 on page 69, listing items for the model protocol. Most significantly, the section on furosemide and the dosage for aminophylline were omitted. The corrected contents for the figure, with underlined changes, are shown below.

1. General Supportive Care
  - ECG monitoring
  - Venous access (IV D5W, t.k.o. or reseat)
  - History, physical assessment
2. Patient Positioning - Upright
3. High Flow O<sub>2</sub>
4. Positive Pressure Ventilation
5. Intubation, if any of the following:
  - Decreased level of consciousness
  - Decreased ventilation (RR>36 or <10)
  - Hypotensive
6. PEEP (intubated cases)
  - 5-10 cm. H<sub>2</sub>O
7. Nitrates
  - 2.5 mg. isosorbide or 0.4 mg. nitroglycerin, sublingual
8. Morphine
  - 2.5 mg. increments @ 5 min. intervals until diastolic 70-100 or systolic 120-160
9. Furosemide
  - 40 mg. (80 mg. if on oral furosemide); Repeat at double dose, if needed
10. Aminophylline
  - 250 mg. in 50 ml. over 30 min.  
(if not on oral theophylline products)

Errata normally appear in a subsequent issue of the Journal. However, the significance of these errors prompted our decision to publish this special errata. We apologize for the omission and any confusion it may have caused. Despite the very limited time and resources under which we produce the Journal, we make no excuses for such errors - We take our responsibility for the content of this publication very seriously. We appreciate the support of our readers and assure that as we grow, we will continue to improve the composition and academic quality of its contents.

The Editorial Staff